



## FEATURES & BENEFITS

- Highly effective 1% iodine
- Formulated with IoPlex, AgroChem's proprietary process that provides a steady, stable supply of free, available iodine.
- Rich 12% emollient package that includes aloe vera, glycerine, lanolin and propylene glycol.
- Forms a physical barrier on teat skin to block pathogens.



## ORDERING INFORMATION

<u>Premier Barrier</u>	
3371B	5 gal
3372B	15 gal
3373B	55 gal
3374B	265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuses, mishandling or application not specifically described on the label.



# PREMIER BARRIER

## SPECIFICATION DATA

<b>Primary Germicide</b>	1% Iodine
<b>pH</b>	4.5 - 5.5, 100%
<b>Color</b>	Dark Brown, Reddish
<b>Fragrance</b>	Iodine
<b>Mix Rate</b>	Ready-To-Use (RTU)
<b>Usage</b>	Post
<b>Emollient Package</b>	12% Total Aloe Vera Glycerine Lanolin Propylene Glycol

## DIRECTIONS FOR USE

**NOTE:** Product is in a Ready-To-Use (RTU) concentration. Do not mix product with water, another teat dip, or other product.

### GENERAL USE

#### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

#### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

## SAFETY & DANGER

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

### FIRST AID / RESPONSE:

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/container to an approved waste disposal plant



26 Freedom Way, Saratoga Springs, NY 12866  
(518) 226-4850 | www.agrochemusa.com