



FEATURES & BENEFITS

- Forms a physical barrier on teat skin to physically block pathogens.
- Dual-germicide system provides quick kill and long-lasting protection.
- Barrier film is easily removed with standard pre-dips.
- Rich skin conditioning and emollient system.



ORDERING INFORMATION

XT Barrier – Part A

3592A 5 gal
3593A 15 gal
3594A 55 gal
3595A 265 gal

XT Barrier– Part B

3582B 5 gal
3583B 15 gal
3584B 55 gal
3585B 265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or application not specifically described on the label.



SPECIFICATION DATA WHEN PART A & PART B ARE MIXED

Primary Germicide	Chlorine Dioxide
pH	3.0, 100%
Color	Green
Fragrance	Chlorine
Mix Rate	1:1
PPM	1,500 - 1,900
Usage	PPost
Emollient Package	10.0% Total Glycerin Lanolin

DIRECTIONS FOR USE

XT BARRIER is a two-part Chlorine Dioxide teat dip and can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

DIRECTIONS FOR USE

Mix Directions: Mix ratio is 1:1. Add equal parts of XT Barrier - PART A and PART B in a separate container. Inverting the container approximately 5-7 times will adequately mix the two solutions. This mixed solution will remain stable for 7 days. After 7 days, discard solution and mix a new solution.

IMPORTANT: Do not add solution back into the original containers for XT Barrier Base or Activator. Product activation could occur. Do not add XT Barrier ready-to-use solution to any other teat dip or product. If transferred from this container to any other, make sure the other container is thoroughly pre-cleaned and bears the proper container labeling for XT Barrier. Use XT Barrier on each teat as an aid in a complete cow-care program to help reduce the spread of organisms which may cause mastitis. The barrier properties of XT Barrier will provide protection until the next milking.

General Use

Teat Sprays can also be foamed or sprayed using recommended equipment. If using a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

SAFETY & DANGER

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant



26 Freedom Way, Saratoga Springs, NY 12866
(518) 226-4850 | www.agrochemusa.com