

FEATURES & BENEFITS

- Resistant to breakdown and loss of activity caused by organic matter and soil – unlike iodine and peroxide-based teat dips.
- Produces powerful germicides once activated, including chlorine dioxide and chlorous acid.
- Will not change pH of healthy skin.
- Delivers a broad-spectrum kill rate in as little as 15 to 20 seconds.
- Contains lactic acid, an Alpha Hydroxy Acid (AHA) which softens and exfoliates rough teat ends to allow for easy removal of damaged skin.



ORDERING INFORMATION

| Xtra-7 - | - Part A | <u> Xtra-7 -</u> | - Part B |
|----------|----------|------------------|----------|
| 36015 | 5 gal | 3651S | 5 gal |
| 3602S | 15 gal | 3652S | 15 gal |
| 3603S | 55 gal | 3653S | 55 gal |
| 36045 | 265 gal | 3654S | 265 gal |



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuses, mishandling or application not specifically described on the label.



26 Freedom Way, Saratoga Springs, NY 12866 (518) 226-4850 | www.agrochemusa.com



SPECIFICATION DATA WHEN PART A & PART B ARE MIXED

| Primary Germicide | Chlorine Dioxide | |
|-------------------|--|--|
| рН | 2.1, 100% | |
| Color | Yellow | |
| Fragrance | Slight, Chlorine | |
| Mix Rate | 1:1 | |
| РРМ | 2,200 - 2,400 | |
| Usage | Pre & Post | |
| Emollient Package | 7.0 % Glycerin Lanolin Propylene Glycol | |
| Pump | 8502 GG.R | |

DIRECTIONS FOR USE

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA-7 after 24 hours.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

SAFETY & DANGER

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant