



## FEATURES & BENEFITS

- Highly effective 0.5% iodine, 2% emollient package with 21.7 ppm of Free Iodine.
- Formulated using AgroChem's proprietary IoPlex method for quick kill, long shelf life and gentler effect on teat skin.
- Withstands freeze/thaw cycles without the loss or fallout of iodine.
- Provide superior softening and conditioning



## ORDERING INFORMATION

### BovaPro 52 Dip & Spray

3251S 5 gal  
 3252S 15 gal  
 3253S 55 gal  
 3254S 265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or application not specifically described on the label.

# BovaPro<sup>52</sup>

## SPECIFICATION DATA

<b>Primary Germicide</b>	0.5% Iodine
<b>pH</b>	4.5 - 5.5, 100%
<b>Color</b>	Dark Brown, Reddish
<b>Fragrance</b>	Iodine
<b>Mix Rate</b>	Ready-To-Use (RTU)
<b>Free Iodine</b>	21.7 ppm
<b>Usage</b>	Pre & Post
<b>Emollient Package</b>	2% Total Glycerine

## DIRECTIONS FOR USE

**NOTE:** Product is in a Ready-To-Use (RTU) concentration. Do not mix product with water, another teat dips, or other product.

### GENERAL USE

#### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

#### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

## SAFETY & DANGER

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

### FIRST AID / RESPONSE:

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/container to an approved waste disposal plant



26 Freedom Way, Saratoga Springs, NY 12866  
 (518) 226-4850 | www.agrochemusa.com