

## **FEATURES & BENEFITS**

- Highly effective 1% iodine, 4% emollient package.
- Blocks pathogens by forming a physical and chemical barrier on teat skin.
- Dries quickly, making BovaPro Barrier Film ideal for dairies bedding on recycled manure solids.
- Very low drip rate.
- Easily removed with standard pre-dips



### **ORDERING INFORMATION**

<u>BovaPro Barrier Film</u>		
3101FM	5 gal	
3102FM	15 gal	
3103FM	55 gal	
3104FM	265 gal	



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misues, mishandling or application not specifically described on the label.



26 Freedom Way, Saratoga Springs, NY 12866 (518) 226-4850 | www.agrochemusa.com

# BOVA Pro BARRIER FILM

# **SPECIFICATION DATA**

Primary Germicide	1% lodine
рН	2.75 - 3.5, 100%
Color	Dark Brown, Reddish
Fragrance	lodine
Mix Rate	Ready-To-Use (RTU)
Usage	Pre
Emollient Package	4% Total Glycerine

# **DIRECTIONS** FOR USE

**NOTE:** Product is in a Ready-To-Use (RTU) concentration. Do not mix product wiht water, an other teat dips, or other product.

#### **GENERAL USE**

#### **Pre Dipping Use**

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### **Post Dipping Use**

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

#### **Extended** Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

## **SAFETY & DANGER**

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/container to an approved waste disposal plant