



FEATURES & BENEFITS

- Highly effective 1% iodine,
- Formulated with IoPlex, AgroChem's proprietary process that provides a steady, stable supply of free, available iodine.
- 10% emollient package that includes glycerine.
- Provides an excellent water resistant, low drop physical barrier on the teat.



ORDERING INFORMATION

BovaGuard Barrier Dip

3171B	5 gal
3172B	15 gal
3173B	55 gal
3174B	265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuses, mishandling or application not specifically described on the label.



26 Freedom Way, Saratoga Springs, NY 12866
(518) 226-4850 | www.agrochemusa.com



SPECIFICATION DATA

Primary Germicide	1% Iodine
pH	4.5 - 5.5, 100%
Color	Dark Brown, Reddish
Fragrance	Iodine
Mix Rate	Ready-To-Use (RTU)
Usage	Post
Emollient Package	12% Total Aloe Vera Glycerine Lanolin Propylene Glycol

DIRECTIONS FOR USE

NOTE: Product is in a Ready-To-Use (RTU) concentration. Do not mix product with water, an other teat dips, or other product.

GENERAL USE

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

SAFETY & DANGER

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/container to an approved waste disposal plant