



## FEATURES & BENEFITS

- Fast-acting chlorhexidine gluconate-based teat dip with a distinctive blue color.
- Extremely effective microbicide active against most gram-positive and gram-negative organisms, certain anaerobes, aerobes and yeasts.
- Available in 0.5% and 1.0% chlorhexidine concentrations.
- Helps decrease intramammary infections due to Strep. agalactiae and Staph. aureus.
- Adheres well, and stays active for up to 5-6 hours longer than other antimicrobial agents.
- Glycerin emollient for superior skin conditioning.



## ORDERING INFORMATION

### Chlordine 110

3401S 5 gal  
3402S 15 gal  
3403S 555 gal  
3404S 265 gal

### Chlordine 55

3431S 5 gal  
3432S 15 gal  
3433S 555 gal  
3434S 265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuses, mishandling or application not specifically described on the label.



26 Freedom Way, Saratoga Springs, NY 12866  
(518) 226-4850 | www.agrochemusa.com

# CHLORDINE

## SPECIFICATION DATA

<b>Primary Germicide</b>	Chlorhexidine Gluconate
<b>pH</b>	6, 100%
<b>Color</b>	Blue
<b>Fragrance</b>	Faint
<b>Mix Rate</b>	Ready-To-Use
<b>Usage</b>	Pre & Post
<b>Emollient Package</b>	Glycerin

## DIRECTIONS FOR USE

### General Use

RTU does not require dilution and can be used immediately.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

### Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

## SAFETY & DANGER

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

**FIRST AID / RESPONSE:**

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/ container to an approved waste disposal plant