



FEATURES & BENEFITS

- Highly concentrated formula provides a cost effective pre and post-dip solution
- Rich skin conditioning and emollient system with lanolin and other emollients.
- Dual germicide delivers a broad spectrum kill rate in little as 15 - 20 seconds.
- Lactic acid provides gentle exfoliation of dead skin and removes keratin build-up known as hyperkeratosis.
- One product for pre and post dipping



ORDERING INFORMATION

XT 22C- Part A

3261A 5 gal
 3262A 15 gal
 3263A 55 gal
 3264A 265 gal

XT 22C Part B

3391B 5 gal
 3392B 15 gal
 3393B 55 gal
 3394B 265 gal



Manufacturer or seller makes no warranty, express or implied, concerning use of the product other than for the purpose indicated on the label. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or application not specifically described on the label.



SPECIFICATION DATA WHEN PART A & PART B ARE MIXED

Primary Germicide	Chlorine Dioxide
pH	6.0, 100%
Color	Green
Fragrance	Chlorine
Mix Rate	1:1:20
PPM	1,000 - 1,200
Usage	Pre & Post
Emollient Package	2.5% Total* Lanolin Emollients <small>*Additional can be added with Aurora blending system</small>
Pump	8502 GG.R or Aurora

DIRECTIONS FOR USE

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for multiple days.

Pre Dipping Use

Perform fore stripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, the full length of each teat with the product.

Allow product to stand for 30 seconds. Wipe teats dry after application, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip, foam, or spray, the full length of each teat. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

SAFETY & DANGER

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant



26 Freedom Way, Saratoga Springs, NY 12866
 (518) 226-4850 | www.agrochemusa.com